

Product datasheet for **TP724301**

Human ADAM15 Protein, hFc Tag

Product data:

Product Type:	Recombinant Proteins
Description:	Human ADAM15 Protein, hFc Tag
Expression Host:	HEK293
Tag:	C-Human Fc
Predicted MW:	The protein has a predicted molecular mass of 100.0 kDa after removal of the signal peptide. The apparent molecular mass of ADAM15-hFc is approximately 100-130 kDa due to glycosylation.
Purity:	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Reconstitution Method:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization.
Storage:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Stability:	12 months from date of despatch
Synonyms:	ADAM 15, MDC-15
Summary:	The protein encoded by this gene is a member of the ADAM (a disintegrin and metalloproteinase) protein family. ADAM family members are type I transmembrane glycoproteins known to be involved in cell adhesion and proteolytic ectodomain processing of cytokines and adhesion molecules. This protein contains multiple functional domains including a zinc-binding metalloprotease domain, a disintegrin-like domain, as well as a EGF-like domain. Through its disintegrin-like domain, this protein specifically interacts with the integrin beta chain, beta 3. It also interacts with Src family protein-tyrosine kinases in a phosphorylation-dependent manner, suggesting that this protein may function in cell-cell adhesion as well as in cellular signaling. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]



[View online »](#)