

Product datasheet for **TP724365**

Human XAGE1A Protein, hFc Tag

Product data:

| | |
|-------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | Human XAGE1A Protein, hFc Tag |
| Expression Host: | HEK293 |
| Tag: | N-Human Fc |
| Predicted MW: | The protein has a predicted molecular mass of 35.2 kDa after removal of the signal peptide. The apparent molecular mass of hFc-XAGE1A is approximately 35-55 kDa due to glycosylation. |
| Purity: | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Reconstitution Method: | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. |
| Storage: | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Stability: | 12 months from date of despatch |
| Summary: | This gene is a member of the XAGE subfamily, which belongs to the GAGE family. The GAGE genes are expressed in a variety of tumors and in some fetal and reproductive tissues. This gene is strongly expressed in Ewing's sarcoma, alveolar rhabdomyosarcoma and normal testis. The protein encoded by this gene contains a nuclear localization signal and shares a sequence similarity with other GAGE/PAGE proteins. Because of the expression pattern and the sequence similarity, this protein also belongs to a family of CT (cancer-testis) antigens. Alternative splicing of this gene, in addition to alternative transcription start sites, results in multiple transcript variants. [provided by RefSeq, Jan 2010] |



[View online »](#)