

Product datasheet for TP724733

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Mouse CD70 (C-mFc)

Species: Mouse

Expression cDNA Clone

or AA Sequence:

Gln39-Pro193

Tag: C-Mouse Fc

Buffer: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Note: Recombinant Mouse CD70 (C-mFc) is produced by Human Cells. The target gene encoding

Gln39-Pro193 is expressed with a C-mFc tag.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-5 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: 12 months from date of despatch

Synonyms: CD70,CD27LG,TNFSF7,TNFSF7G,CD27L

Summary: Cluster of Differentiation 70 (CD70) is also known as CD27 ligand (CD27L / CD27LG), TNFSF7,

TNFSF7G, is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and is a surface antigen found on activated T-and B-lymphocytes and mature dendritic cells. Binding of CD70 to its receptor CD27 induces in priming, effector functions, differentiation and memory formation of T-cells, and thus is invloved in the biological processes including T-cell activation, the proliferation of costimulated T-cells, as well as the generation of cytolytic T-cells. CD70 on T cells provides costimulatory signals that are required for T cell proliferation, clonal expansion and the promotion of effector T cell formation. CD70 on mouse B cell has been shown to inhibit terminal differentiation of activated B cells into plasma cells and enhances commitment to memory B cell responses. CD70 induces proliferation and IFNÎ³

production, on NK cells.

