

## Product datasheet for **TP726583**

### **Tnfrsf13c Mouse Recombinant Protein**

#### **Product data:**

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant Mouse BAFFR (C-Fc)
<b>Species:</b>	Mouse
<b>Expression cDNA Clone or AA Sequence:</b>	Ser10-Ala 71
<b>Tag:</b>	C-Fc
<b>Buffer:</b>	Lyophilized from a 0.2 um filtered solution of PBS, pH7.4.
<b>Note:</b>	Recombinant Mouse Tumor Necrosis Factor Receptor Superfamily Member 13C is produced by our Mammalian expression system and the target gene encoding Ser10-Ala 71 is expressed with a Fc tag at the C-terminus.
<b>Stability:</b>	12 months from date of despatch
<b>Locus ID:</b>	72049
<b>UniProt ID:</b>	<a href="#">Q9D8D0</a>
<b>Summary:</b>	Tumor necrosis factor receptor superfamily, member 13C (TNFRSF13C) also known as B-cell-activating factor receptor (BAFFR) and CD268 antigen, is a member of the tumor necrosis factor receptor superfamily. BAFF promotes the survival of B cells and is essential for B cell maturation. BAFF binds to three TNF receptor superfamily members: B-cell maturation antigen (BCMA/TNFRSF17), transmembrane activator and calcium-modulator and cyclophilin ligand interactor (TACI/TNFRSF13B) and BAFF receptor (BAFF R/BR3/TNFRSF13C). These receptors are type III transmembrane proteins that lack a signal peptide. BAFF R is highly expressed in spleen, lymph node and resting B cells. It is also expressed at lower levels in activated B cell, in resting CD4+ T cells, in thymus and peripheral blood leukocytes. BAFF knockout mice lack mature B cells. Similarly, A/WySnJ mice that are defective in BAFF-R intracellular signaling also lack mature B cells, suggesting that BAFF R is the critical receptor for BAFF during B lymphopoiesis. It has been proposed that abnormally high levels of BAFFR/TNFRSF13C (CD268) may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells.


[View online »](#)