

Product datasheet for **TP726657**

Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Human ALK-2 (C-Fc)
Species:	Human
Expression cDNA Clone or AA Sequence:	Asp23 ^Δ Val124
Tag:	C-Fc
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS,pH7.4.
Note:	Recombinant Human Receptor Protein Serine/Threonine Kinase is produced by our Mammalian expression system and the target gene encoding Asp23 ^Δ Val124 is expressed with a Fc tag at the C-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Synonyms:	activin A receptor, type I; Activin RIA; ActivinRIA; ACTRI; ACVR1; ACVR1A; ACVRLK2; ALK2; ALK-2; FOP; SKR1; TSRI
Summary:	Activin RIA, also known as ALK-2, TSK-7L, SKR1, TSR-I, and ACTR-I, is a glycosylated 65 kDa type I receptor in the TGF-beta serine/threonine kinase receptor family. Binding of TGF-beta superfamily ligands induces formation of a heterotetrameric complex that contains two chains each of a type I and a type II receptor in multiple combinations. The type II receptors phosphorylate the type I receptors which then phosphorylate and activate Smad signal transduction proteins. Activin RIA functions in a wide variety of growth and differentiation processes including gastrulation, skeletal system development, and cardiac morphogenesis. BMP signaling through Activin RIA is enhanced by the direct interaction between Activin RIA and RGM-B/DRAGON, a BMP coreceptor that also associates with other type I and type II receptors. Activin RIA can additionally phosphorylate the coreceptor Endoglin and is required for the inhibitory effect of Endoglin on prostate cancer cell motility.



[View online »](#)