

Product datasheet for **TP726784**

IL17re Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Mouse IL-17RE (C-6His)
Species:	Mouse
Expression cDNA Clone or AA Sequence:	Ala115His414
Tag:	C-6His
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS,pH7.4.
Note:	Recombinant Mouse Interleukin-17 Receptor E is produced by our Mammalian expression system and the target gene encoding Ala115His414 is expressed with a 6His tag at the C-terminus.
Storage:	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at $2-8^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Stability:	12 months from date of despatch
Locus ID:	57890
UniProt ID:	Q8BH06
Synonyms:	IL-17 RE; IL-17 receptor E; IL17RE; IL-17RE; interleukin 17 receptor E

[View online »](#)

Summary:

Interleukin 17 Receptor E (IL 17 RE) is an approximately 70 kDa (predicted) transmembrane protein in the family of IL 17 receptors. IL 17 RE is expressed on keratinocytes, mucosal epithelial cells, Th17 cells, and gamma δ T cells. It associates with the widely expressed IL 17 RA to form a heterodimeric receptor for IL-17C. IL-17C binds to IL 17 RE with high affinity and to IL 17 RA with low affinity. IL 17C expression is induced by inflammatory stimulation in colon and airway epithelial cells, keratinocytes, CD4+ T cells, macrophages, and dendritic cells. It is up regulated in various chronic inflammatory diseases including psoriasis, cystic fibrosis, and chronic obstructive pulmonary disease (COPD). IL 17 RE is reciprocally down regulated in psoriatic lesions. The interaction of IL 17C with IL 17 RE promotes mucosal immunity through the induction of anti bacterial peptides and pro inflammatory cytokines and chemokines. IL 17C action supports the integrity of the colon epithelium following infection induced damage but also contributes to psoriatic skin thickening and the progression of arthritis. IL 17C is additionally up regulated in Th17 cell dependent autoimmunity. In this setting, it exacerbates disease severity by inducing Th17 cell production of IL 17A, IL 17F, IL 22, CCR6, and CCL20. The up regulation of IL 17 RE in hepatocellular carcinoma is associated with poor prognosis.