

## Product datasheet for **TP726792**

### Human Recombinant Protein

#### Product data:

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant Human CD200 Receptor 1/CD200R1 (C-6His)
<b>Species:</b>	Human
<b>Expression cDNA Clone or AA Sequence:</b>	Ala27-Leu266
<b>Tag:</b>	C-His
<b>Buffer:</b>	Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
<b>Note:</b>	Recombinant Human CD200 Receptor 1 is produced by our Mammalian expression system and the target gene encoding Ala27-Leu266 is expressed with a 6His tag at the C-terminus.
<b>Storage:</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Stability:</b>	12 months from date of despatch
<b>Synonyms:</b>	Cell Surface Glycoprotein CD200 Receptor 1; CD200 Cell Surface Glycoprotein Receptor; Cell Surface Glycoprotein OX2 Receptor 1; CD200R1; CD200R; CRTR2; MOX2R; OX2R
<b>Summary:</b>	Cell surface glycoprotein CD200 Receptor 1 (CD200R1) is the receptor for the CD200 (OX-2) membrane glycoprotein. CD200R1 contains one C2- type Ig-like domain and one V-type Ig-like domain within its extracellular domain and a PTB-signaling motif in cytoplasmic domain. CD200R1 and CD200 associate via their respective N-terminal Ig-like domains. CD200R1 is restricted primarily to mast cells, basophils, macrophages, and dendritic cells. It propagates inhibitory signals despite its lacking a cytoplasmic ITIM (immunoreceptor tyrosinebased inhibitory motif). The receptor-substrate interaction may function as a myeloid downregulatory signal.



[View online »](#)