

Product datasheet for TP726861

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Human FAD-linked Sulfhydryl Oxidase ALR/GFER (N-6His)

Species: Human

Expression cDNA Clone

or AA Sequence:

Met1-Asp125

Tag: N-His

Buffer: Lyophilized from a 0.2 um filtered solution of 4mM HCl.

Note: Recombinant Human Growth Factor, Augmenter of Liver Regeneration is produced by our

E.coli expression system and the target gene encoding Met1-Asp125 is expressed with a 6His

tag at the N-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: 12 months from date of despatch

Synonyms: FAD-linked sulfhydryl oxidase ALR;GFER;Augmenter of liver

regeneration;hERV1;Hepatopoietin;GFER;ALR;HERV1;HPO

Summary: GFER is a hepatotrophic growth factor and flavin-linked sulfhydryl oxidase which belongs to

the Erv1/ALR family of proteins. GFER is widely expressed in various human tissues. They are two isoforms of this protein. Isoform 1 could regenerate the redox-active disulfide bonds in CHCHD4/MIA40, a chaperone essential for disulfide bond formation and protein folding in the mitochondrial intermembrane space. The reduced form of CHCHD4/MIA40 forms a transient intermolecular disulfide bridge with GFER/ERV1, resulting in regeneration of the essential disulfide bonds in CHCHD4/MIA40, while GFER/ERV1 becomes re-oxidized by donating electrons to cytochrome c or molecular oxygen. Isoform 2 may act as an autocrine hepatotrophic growth factor promoting liver regeneration. GFER could also induce the expression of S-adenosylmethionine decarboxyl-ase and ornithine decarboxylases (ODC). S-adenosylmethionine decarboxyl-ase and ornithine decarboxylases play an important role in

the synthesis of polyamines.

