

## **Product datasheet for TP726985**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ERO1L (ERO1A) Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Human ERO1-Like Protein α/ERO1L (C-6His)

Species: Human

**Expression cDNA Clone** 

or AA Sequence:

Glu24-His468

Tag: C-His

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

**Note:** Recombinant Human ERO1-Like Protein alpha is produced by our Mammalian expression

system and the target gene encoding Glu24-His468 is expressed with a 6His tag at the C-

terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Stability:** 12 months from date of despatch

Locus ID: 30001 UniProt ID: Q96HE7

Synonyms: ERO1-Like Protein Alpha; ERO1-L; ERO1-L-Alpha; Endoplasmic Oxidoreductin-1-Like Protein;

Oxidoreductin-1-L-Alpha; ERO1L

**Summary:** ERO1-Like Protein α (ERO1L) is an enzyme that belongs to the EROs family. ERO1L is

expressed at high level in esophagus and upper digestive tract. ERO1L is an essential oxidoreductase that oxidizes proteins in the endoplasmic reticulum to produce disulfide bonds. ERO1L acts by oxidizing directly P4HB/PDI isomerase through a direct disulfide exchange. It associates with ERP44, demonstrating that it does not oxidize all PDI related proteins and can discriminate between PDI and related proteins. Its reoxidation probably involves electron transfer to molecular oxygen via FAD. ERO1L may be responsible for a significant proportion of reactive oxygen species (ROS) in the cell. ERO1L responses to temperature stimulus, protein thiol-disulfide exchange, protein folding with or without

chaperone cofactor and transport.

**Protein Pathways:** Vibrio cholerae infection

