

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP727045

FGFR3 Human Recombinant Protein

Product data:

Recombinant Proteins
Recombinant Human Fibroblast Growth Factor Receptor 3/FGF R3 (C-Fc)
Human
Glu23-Gly375
C-Fc
Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Recombinant Human Fibroblast Growth Factor Receptor 3 is produced by our Mammalian expression system and the target gene encoding Glu23-Gly375 is expressed with a Fc tag at the C-terminus.
Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
12 months from date of despatch
2261
<u>P22607</u>
Fibroblast growth factor receptor 3; FGFR-3; CD333; FGFR3; JTK4; IIIc



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE FGFR3 Human Recombinant Protein – TP727045

cytoskeleton

Summary:	Fibroblast growth factors (FGFs) are involved in a multitude of physiological and pathological cellular processes. The biological activities of the FGFs are mediated by a family of type I transmembrane tyrosine kinases which undergo dimerization and autophosphorylation after ligand binding. Four distinct genes encoding closely related FGF receptors, FGF R1-4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. Multiple forms of FGF R1-3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGF R1 and 2 results in receptors containing all three Ig domains, referred to as the $\hat{1}\pm$ isoform, or only IgII and IgIII, referred to as the $\hat{1}^2$ isoform. Only the $\hat{1}\pm$ isoform has been identified for FGF R3 and FGF R4. Additional splicing events for FGF R1-3, involving the C-terminal half of the IgIII domain encoded by two mutually exclusive alternative exons, generate FGF receptors with alternative IgIII domains (IIIb and IIIc). The complex patterns of expression of these receptors as well as the specificity of their interactions with the various FGF Iigand family members are under investigation.
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Bladder cancer, Endocytosis, MAPK signaling pathway, Pathways in cancer, Regulation of actin