

## **Product datasheet for TP727086**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Il5ra Mouse Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Mouse Interleukin-5 Receptor Subunit Alpha/IL-5 Rα (C-6His)

Species: Mouse

**Expression cDNA Clone** 

or AA Sequence:

Asp18-His339

Tag: C-His

**Buffer:** Lyophilized from a 0.2 um filtered solution of PBS, 5% Trehalose, pH 7.4.

**Note:** Recombinant Mouse Interleukin-5 Receptor Subunit Alpha is produced by our Mammalian

expression system and the target gene encoding Asp18-His339 is expressed with a 6His tag at

the C-terminus.

**Stability:** 12 months from date of despatch

Locus ID: 16192 UniProt ID: <u>P21183</u>

Summary: Interleukin 5 Receptor alpha (IL-5 Rα), also known as CD125, is a hematopoietin receptor that

plays a dominant role in eosinophil biology. Mature mouse IL-5 Rα consists of a 322 amino acid (aa) extracellular domain (ECD) with a WSxWS motif and a four cysteine motif, a 22 aa transmembrane segment, and a 54 aa cytoplasmic domain. The high affinity receptor for IL-5 is a complex that consists of the ligand binding IL-5 Rα and the transmembrane common β chain (βc/CD131) which is shared with the receptor complexes for IL-3 and GM-CSF. IL-5 Rα binds IL-5 at low affinity and then associates with preformed βc oligomers to form the signaling-competent receptor complex. IL-5 stimulation of CD34+ hematopoietic progenitor cells induces the up-regulation of transmembrane IL-5 Rα followed by eosinophilic

differentiation and activation. IL-5 Rα also promotes the differentiation of basophils and B cells. Exposure of mature eosinophils to IL-5 attenuates their IL-5 responsiveness by inducing

the down-regulation of surface IL-5 Rα and increased production of soluble IL-5 Rα.

