

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP727098

Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Human Receptor-Type Tyrosine-Protein Kinase FLT3/FLT3 (C-Fc)
Species:	Human
Expression cDNA Clone or AA Sequence:	Asn27-Asn541
Tag:	C-Fc
Buffer:	Supplied as a 0.2 um filtered solution of PBS, pH 7.4.
Note:	Recombinant Human Receptor-Type Tyrosine-Protein Kinase FLT3/FLT3 is produced by our Mammalian expression system and the target gene encoding Asn27-Asn541 is expressed with a Fc tag at the C-terminus.
Storage:	Store at < -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.
Stability:	12 months from date of despatch
Synonyms:	Receptor-Type Tyrosine-Protein Kinase FLT3; FL Cytokine Receptor; Fetal Liver Kinase-2; FLK-2; Fms-Like Tyrosine Kinase 3; FLT-3; Stem Cell Tyrosine Kinase 1; STK-1; CD135; FLT3; FLK2; STK1
Summary:	The Flt-3 (fms-like tyrosine kinase) receptor, also named Flk-2and Stk-1is a member of the class III subfamily of receptor tyrosine kinases that also includes KIT, the receptor for SCF and FMS, the receptor for M-CSF. The extracellular region of these receptors contains five immunoglobulin-like domains and the intracellular region contains a split kinase domain. Human Flt-3 cDNA encodes a 993 amino acid (aa) residue type I membrane protein with a 26 aa residue signal peptide, a 515 aa extracellular domain with 10 potential N-linked glycosylation sites, a 21 aa residue transmembrane domain and a 431 aa residue cytoplasmic domain. Flt-3 expression has been detected in various tissues, including placenta, gonads, and tissues of nervous and hematopoietic origin. Among hematopoietic cells, the expression of Flt-3 was found to be restricted to the highly enriched stem/progenitor cell populations. The ligand for Flt-3 (FL) has been identified to be a transmembrane protein with structural homology to M-CSF and SCF. Recombinant soluble Flt-3/Fc chimeric protein has been shown to bind FL with high affinity and is a potent FL antagonist.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US