

## **Product datasheet for TP727211**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Human Receptor Tyrosine Kinase MerTK/MERTK/MER (C-6His)

Species: Human

**Expression cDNA Clone** 

or AA Sequence:

Met177-Ala499

Tag: C-His

**Buffer:** Lyophilized from a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.

**Note:** Recombinant Human Tyrosine-protein kinase Mer is produced by our Mammalian expression

system and the target gene encoding Met1-Ala323 is expressed with a 6His tag at the C-

terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Stability:** 12 months from date of despatch

Synonyms: Tyrosine-protein kinase Mer/Proto-oncogene c-Mer/Receptor tyrosine kinase

MerTK/MERTK/MER

**Summary:** Tyrosine-protein kinase Mer (MERTK) is a single-pass type I membrane protein which belongs

to the MER/AXL/TYRO3 receptor kinase family. MERTK include two fibronectin type-III domains, two Ig-like C2-type domains, and one tyrosine kinase domain. It can't be expressed in normal B- and T-lymphocytes, but it is usually expressed in numerous

neoplastic B- and T-cell lines. MERTK could regulate many physiological processes, such as cell survival, migration, differentiation. It was demonstrated that the MERTK plays critical role

in the engulfment and efficient clearance of apoptotic cells, platelet aggregation, and cytoskeleton reorganization. Not only these, it also plays an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3. In addition, MERTK could regulate rod outer segments fragments phagocytosis in the retinal pigment epithelium (RPE), deficiency in MERTK are the cause of retinitis pigmentosa.

