

Product datasheet for TP727289

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL2 Receptor beta (IL2RB) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Human IL-2 Receptor Subunit β/IL-2RB/CD122 (C-Fc)

Species: Human

Expression cDNA Clone

or AA Sequence:

Ala27-Asp239

Tag: C-Fc

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Note: Recombinant Human Interleukin-2 Receptor subunit beta is produced by our Mammalian

expression system and the target gene encoding Ala27-Asp239 is expressed with a Fc tag at

the C-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: 12 months from date of despatch

Locus ID: 3560 **UniProt ID:** P14784

Synonyms: Interleukin-2 receptor subunit beta;IL2RB;IL-2 receptor subunit beta;IL-2R subunit beta;High

affinity IL-2 receptor subunit beta;CD122

Summary: Human IL-2RB, also known asinterleukin-2 receptor subunit beta,is the receptor for

interleukin-2. IL2 receptor complex is involved in receptor mediated endocytosis and

transduces the mitogenic signals of IL2. IL2 receptor complex has three forms with respect to ability to bind IL2. IL-2RB is belonged to a type I membrane protein, and has a 26 residue signal peptide, a 214 residue extracellular region, a 25 residue transmembrane region and a 286 residue cytoplasmic domain. IL-2RB is the subunit critical for receptor-mediated signaling via physically or functionally coupling to other signaling molecules, such as the Jak-STAT and

Src-family protein tyrosine kinase although it lacks apparent catalytic motifs.

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Endocytosis, Jak-STAT signaling pathway

