

Product datasheet for TP727290

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Lgmn Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Mouse Legumain/Asparaginyl Endopeptidase (C-6His)

Species: Mouse

Expression cDNA Clone

or AA Sequence:

Val18-Tyr435

Tag: C-His

Buffer: Supplied as a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, 20% Glycerol, pH 8.0.

Note: Recombinant Mouse Legumain/Asparaginyl Endopeptidase is produced by our Mammalian

expression system and the target gene encoding Val18-Tyr435 is expressed with a 6His tag at

the C-terminus.

Storage: Store at < -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.

Stability: 12 months from date of despatch

Locus ID: 19141 **UniProt ID:** 089017

Synonyms: Legumain;Lgmn;Asparaginyl endopeptidase;Protease cysteine 1;Prsc1

Summary: Mouse Legumain, also known as LGMN, is a cysteine protease belonging to peptidase family

C13 and is expressed in kidney and placenta abundantly. LGMN has a strict specificity for hydrolysis of asparaginyl bonds. It can also cleave aspartyl bonds slowly, especially under acidic conditions. The mammalian legumain is involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system. It plays a role in the regulation of cell proliferation via its role in EGFR degradation. Legumain deficiency causes the accumulation of pro-Cathepsins B, H and L, another group of lysosomal cysteine

proteases. Overexpression of Legumain in tumors is significant for invasion/metastasis. Mammalian legumain is inhibited by iodoacetamide and maleimides. Legumain activation

appears to be autocatalytic and can be triggered by acidic pH.

