

## Product datasheet for **TP727296**

### PD-L1 (Cd274) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Mouse PD-L1/B7-H1/CD274 (C-Fc)
Species:	Mouse
Expression cDNA Clone or AA Sequence:	Phe19-Thr238
Tag:	C-Fc
Buffer:	Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Note:	Recombinant Mouse Programmed Cell Death 1 Ligand 1 is produced by our Mammalian expression system and the target gene encoding Phe19-Thr238 is expressed with a Fc tag at the C-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Locus ID:	60533
UniProt ID:	<u><a href="#">Q9EP73</a></u>
Synonyms:	Programmed cell death 1 ligand 1;Cd274; programmed cell death 1 ligand 1;PD-L1;PDCD1 ligand 1;programmed death ligand 1;B7 homolog 1;B7-H1;CD274



[View online »](#)

**Summary:**

Mouse Programmed cell death 1 ligand 1(Cd274,PD-L1), is a member of the growing B7 family of immune proteins.It involved in the costimulatory signal essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.B7-H1 has been identified as one of two ligands for programmed death1 (PD1), a member of the CD28 family of immunoreceptors. B7-H1 is constitutively expressed in several organs such as heart, skeletal muscle B7-H1 expression is upregulated in a small fraction of activated T and B cells and a much larger fraction of activated monocytes. The costimulatory function of B7-H1 is critical for enhancing maturation and differentiation of T-cells in lymphoid organs.B7-H1 expression is also induced in dendritic cells and keratinocytes after IFN gamma stimulation. Interaction of B7-H1 with PD1 results in inhibition of TCR-mediated proliferation and cytokine production. The B7-H1:PD1 pathway is involved in the negative regulation of some immune responses and may play an important role in the regulation of peripheral tolerance.