

# **Product datasheet for TP727518**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL1 Receptor II (IL1R2) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant Human IL-1 Receptor Type 2/IL-1R-2 (C-6His)

Species: Human

**Expression cDNA Clone** 

or AA Sequence:

Phe14-Glu343

Tag: C-His

**Buffer:** Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

**Note:** Recombinant Human Interleukin-1 receptor type 2/IL-1R-2 is produced by our Mammalian

expression system and the target gene encoding Phe14-Glu343 is expressed with a 6His tag

at the C-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Stability:** 12 months from date of despatch

**Locus ID:** 7850 **UniProt ID:** P27930

Synonyms: Interleukin-1 receptor type 2; IL-1R-2; IL-1RT-2; IL-1RT2; CD121 antigen-like family member B;

CDw121b; IL-1 type II receptor; Interleukin-1 receptor beta; IL-1R-beta; Interleukin-1 receptor

type II; CD121b

Summary: Interleukin-1 receptor type 2 (IL1R2) belongs to the interleukin-1 receptor family. Two distinct

types of IL1 receptors which are able to bind IL1 specifically have been identified, designated as IL1RI (IL1RA) and IL1RII (IL1RB). IL1 receptor type II is a 68 kDa transmembrane protein found on B lymphocytes, neutrophils, monocytes, large granular leukocytes and endothelial cells. IL1R2 is non-signaling receptor forIL1A, IL1B and IL1RN, reduces IL1B activities. IL1R2 serves as a decoy receptor by competetive binding to IL1B and preventing its binding to IL1R1. IL1R2 modulates cellular response through non-signaling association with IL1RAP after binding to IL1B. IL1R2 (membrane and secreted forms) preferentially binds IL1B and poorly IL1A and IL1RN. The secreted IL1R2 recruits secreted IL1RAP with high affinity; this complex formation may be the dominant mechanism for neutralization of IL1B by secreted/soluble

receptors.





### IL1 Receptor II (IL1R2) Human Recombinant Protein – TP727518

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, MAPK signaling pathway