

Product datasheet for TP727525

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eph receptor A2 (EPHA2) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Human Ephrin A Receptor 2/EphA2 (C-Fc)

Species: Human

Expression cDNA Clone

or AA Sequence:

Ala24-Asn534

Tag: C-Fc

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Note: Recombinant Human Ephrin A Receptor 2 is produced by our Mammalian expression system

and the target gene encoding Ala24-Asn534 is expressed with a Fc tag at the C-terminus.

Stability: 12 months from date of despatch

Locus ID: 1969 **UniProt ID:** P29317

Summary: Ephrin type-A receptor 2/EphA2 is a member of the Eph receptor tyrosine kinase family which

binds Ephrins A1, 2, 3, 4, and 5. A and B class Eph proteins have a common structural

organization. Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-

A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is

referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. EphA2 becomes autophosphorolated following ligand

is referred to as reverse signaling. EphA2 becomes autophosphorylated following ligand binding and then interacts with SH2 domain-containing PI3-kinase to activate MAPK pathways. Reverse signaling is also propagated through the Ephrin ligand. Transcription of

EphA2 is dependent on the expression of E-Cadherin, and can be induced by p53 family transcription factors. EphA2 is upregulated in breast, prostate, and colon cancer vascular endothelium. Its ligand, EphrinA1, is expressed by the local tumor cells. In some cases, EphA2 and EphrinA1 are expressed on the same blood vessels. EphA2 signaling cooperates with

VEGF receptor signaling in promoting endothelial cell migration.

