

## **Product datasheet for TP727548**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **MET Human Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Human Hepatocyte Growth Factor Receptor/HGF R/cMet(C-6His)

Species: Human

**Expression cDNA Clone** 

or AA Sequence:

Glu25-Gly519

Tag: C-His

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

**Note:** Recombinant Human Hepatocyte Frowth Factor Receptor is produced by our Mammalian

expression system and the target gene encoding Glu25-Gly519 is expressed with a 6His tag at

the C-terminus.

**Stability:** 12 months from date of despatch

Locus ID: 4233 UniProt ID: P08581

**Summary:** Hepatocyte growth factor receptor (HGF R) is a glycosylated receptor tyrosine kinase that

plays a central role in epithelial morphogenesis and cancer development. HGF R is

synthesized as a single chain precursor which undergoes cotranslational proteolytic cleavage. Mature HGF R is a disulfide-linked dimer composed of a 50 kDa extracellular  $\hat{l}\pm$  chain and a 145 kDa transmembrane  $\hat{l}^2$  chain. Proteolysis and alternate splicing generate additional forms of human HGF R which either lack of the kinase domain, consist of secreted extracellular domains, or are deficient in proteolytic separation of the  $\hat{l}\pm$  and  $\hat{l}^2$  chains. The sema domain, which is formed by both  $\hat{l}\pm$  and  $\hat{l}^2$  chains of HGF R, mediates both ligand binding and receptor dimerization. HGF stimulation induces HGF R downregulation via internalization and

proteasomedependent degradation. Paracrine induction of epithelial cell scattering and branching tubulogenesis results from the stimulation of HGF R on undifferentiated epithelium

by HGF released from neighboring mesenchymal cells.

