

Product datasheet for TP727686

CD23 (FCER2) Human Recombinant Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
Description:	Recombinant Human Fc epsilon RII/CD23 (N-8His)
Species:	Human
Expression cDNA Clone or AA Sequence:	Asp48-Ser321
Tag:	N-His
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Note:	Recombinant Human Low Affinity Fc Epsilon RII is produced by our Mammalian expression system and the target gene encoding Asp48-Ser321 is expressed with a 8His tag at the N-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Locus ID:	2208
UniProt ID:	<u>P06734</u>
Synonyms:	Low affinity immunoglobulin epsilon Fc receptor; BLAST-2; C-type lectin domain family 4 member J; Fc-epsilon-RII; Immunoglobulin E-binding factor; Lymphocyte lgE receptor; CD23; FCER2; CD23A; CLEC4J; FCE2; IGEBF
Summary:	Low affinity immunoglobulin epsilon Fc receptor(CD23) is a secreted and single-pass type II membrane protein which is also exists as a soluble excreted form. There are two forms of CD23: CD23a and CD23b. CD23a is present on follicular B cells, whereas CD23b requires IL-4 to be expressed on T-cells, monocytes, Langerhans cells, eosinophils, and macrophages. Unlike many of the antibody receptors, CD23/FCER2 is a C-type lectin. It is found on mature B cells, activated macrophages, eosinophils, follicular dendritic cells, and platelets. In flow cytometry, CD23/FCER2 is helpful in the differentiation of chronic lymphocytic leukemia (CD23-positive) from mantle cell leukemia (CD23-negative). CD23/FCER2 can also be demonstrated in germinal centre B-cells using immunohistochemistry, but it is not present in the resting cells of the surrounding mantle zone. CD23/FCER2 has essential roles in the regulation of IgE production and in the differentiation of B-cells (it is a B-cell-specificantigen).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD23 (FCER2) Human Recombinant Protein – TP727686

Protein Families: Secreted Protein, Transmembrane

Protein Pathways: Hematopoietic cell lineage

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US