

Product datasheet for **TP727713**

IL5RA Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Human Interleukin-5 Receptor Subunit Alpha/IL-5 R $\hat{\pm}$ (C-6His)
Species:	Human
Expression cDNA Clone or AA Sequence:	Asp21-Glu335
Tag:	C-His
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Note:	Recombinant Human Interleukin-5 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Asp21-Glu335 is expressed with a 6His tag at the C-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Locus ID:	3568
UniProt ID:	<u>Q01344</u>
Synonyms:	Interleukin-5 receptor subunit alpha; IL-5 receptor subunit alpha; IL-5R subunit alpha; IL-5R-alpha; IL-5RA; CDw125; CD125; IL5RA; IL5R
Summary:	Interleukin-5 Receptor alpha (IL-5R $\hat{\pm}$, CD125) is a 60 kDa hematopoietin receptor that plays a dominant role in eosinophil biology. Mature human IL-5 R $\hat{\pm}$ consists of a 322 aa extracellular domain (ECD) with a WSxWS motif and a four cysteine motif, a 20 aa transmembrane segment, and a 58 aa cytoplasmic domain. Within the ECD, human IL-5R $\hat{\pm}$ shares 71% aa sequence identity with mouse and rat IL-5 R $\hat{\pm}$. Alternate splicing of human IL-5 R $\hat{\pm}$ generates soluble secreted forms which function as IL-5 antagonists. The high affinity receptor for IL-5 is a complex that consists of the ligand binding IL-5 R $\hat{\pm}$ and the transmembrane common $\hat{\imath}^2$ chain ($\hat{\imath}^2$ c/CD131) which is shared with the receptor complexes for IL-3 and GM-CSF. IL-5 R $\hat{\pm}$ binds IL-5 at low affinity and then associates with preformed $\hat{\imath}^2$ c oligomers to form the signaling competent receptor complex. IL-5 stimulation of CD34+ hematopoietic progenitor cells induces the up-regulation of transmembrane IL-5R $\hat{\pm}$ followed by eosinophilic differentiation and activation.


[View online »](#)

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling pathway