

## Rocky

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Product datasheet for TP727750

## **Btnl2 Mouse Recombinant Protein**

## **Product data:**

Product Type:	Recombinant Proteins
Description:	Recombinant Mouse Butyrophilin-like Protein 2/BTNL2 (C-6His)
Species:	Mouse
Expression cDNA Clone or AA Sequence:	Asp27-Ser452
Tag:	C-His
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Note:	Recombinant Mouse Butyrophilin-like Protein 2 is produced by our Mammalian expression system and the target gene encoding Asp27-Ser452 is expressed with a 6His tag at the C-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Locus ID:	547431
UniProt ID:	<u>070355</u>
Synonyms:	Butyrophilin-like protein 2; Btnl2; Gm315; Ng9
Summary:	Butyrophilin-like 2 (BTNL2) is a member of the BTN/MOG Ig-superfamily and functions as a negative regulator of immune cell activation. Mouse BTNL2 is type I transmembrane glycoprotein that contains an extracellular domain (ECD), a transmembrane region and a short cytoplasmic domain. The ECD features two V-type Ig-like domains, two C-type Ig-like domains, and four glycosylation sites. BTNL2 is expressed in epithelial cells of the small intestine, colonic dendritic cells, and in cells of the lymph node. BTNL2 expression is upregulated in T cells following activation, a characteristic BTNL2 shares with the homologous B7 family of costimulatory molecules. BTNL2 negatively regulates T cells by inhibiting proliferation and inflammatory cytokine secretion. It also increases the expression of FoxP3 in T cells to promote regulatory T cell development. Single nucleotide polymorphisms in BTNL2 are associated with a risk for sporadic prostate cancer, rheumatoid arthritis, sarcoidosis, ulcerative colitis, and other inflammatory diseases.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US