

Product datasheet for TP727765

Dpt Mouse Recombinant Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Recombinant Proteins |
|--|--|
| Description: | Recombinant Mouse Dermatopontin/DPT (C-Fc) |
| Species: | Mouse |
| Expression cDNA Clone or AA Sequence: | GIn19-Val201 |
| Tag: | C-Fc |
| Buffer: | Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4. |
| Note: | Recombinant Mouse Dermatopontin is produced by our Mammalian expression system and the target gene encoding Gln19-Val201 is expressed with a Fc tag at the C-terminus. |
| Storage: | Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Stability: | 12 months from date of despatch |
| Locus ID: | 56429 |
| UniProt ID: | <u>Q9QZZ6</u> |
| Synonyms: | Early quiescence protein 1; EQ-1; Tyrosine-rich acidic matrix protein; TRAMP |
| Summary: | Dermatopontin is a widely expressed noncollagenous protein component of the extracellular matrix. It is a 22 kDa molecule that is tyrosine sulfated but not glycosylated. Dermatopontin is down regulated in fibrotic growths such as leiomyoma and scar tissue, inhibits cell proliferation, accelerates collagen fibril formation, and stabilizes collagen fibrils against low- temperature dissociation, Dermatopontin deficient mice exhibit altered collagen matrix deposition and organization. Dermatopontin seems to mediate adhesion by cell surface integrin binding, may serve as a communication link between the dermal fibroblast cell surface and its extracellular matrix environment, and enhances TGFB1 activity (By similarity). Dermatopontin promotes bone mineralization under the control of the vitamin D receptor |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

and inhibits BMP-2 effects on osteoblast precursors.