

Product datasheet for TP727768

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Lymphocyte Activation Gene 3 (LAG3) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Human LAG-3/CD233 (C-Fc)

Species: Human

Expression cDNA Clone

or AA Sequence:

Leu23-Leu450

Tag: C-Fc

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Note: Recombinant Human Lymphocyte Activation Gene 3 Protein is produced by our Mammalian

expression system and the target gene encoding Leu23-Leu450 is expressed with a Fc tag at

the C-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: 12 months from date of despatch

Locus ID: 3902 **UniProt ID:** P18627

Synonyms: Lymphocyte activation gene 3 protein;LAG3;LAG-3;Protein FDC;CD223

Summary: Human Lymphocyte activation gene 3 protein(LAG3) is a member of immunoglobulin (Ig)

superfamily. LAG3 contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons.

LAG3 is involved in lymphocyte activation and can bind to HLA class-II antigens. It is

selectively expressed in activated T and NK cells. LAG3 has a negative regulatory function in T cells and acts as as a new marker of T cell induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signaling. It can lead to

increased antigen-specific T-cell responses in vivo. LAG-3 has higher affinity to MHC class II

than CD4.

Protein Families: Transmembrane

