

Product datasheet for TP727810

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd40lg Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant Mouse CD40 ligand/CD40LG (N-6His)

Species: Mouse

Expression cDNA Clone

or AA Sequence:

Met112-Leu260

Tag: N-His

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 200mM NaCl, 0.1mM EDTA, pH 7.0.

Note: Recombinant Mouse CD40 ligand is produced by our Mammalian expression system and the

target gene encoding Met112-Leu260 is expressed with a 6His tag at the N-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: 12 months from date of despatch

Locus ID: 21947 UniProt ID: P27548

Summary: CD40 Ligand, also known as TNFSF5, CD154, is a type II transmembrane glycoprotein

member of the TNF superfamily. Mature mouse CD40 Ligand consists of a 22 amino acid (aa) cytoplasmic domain, a transmembrane segment, and a 214 aa extracellular region. CD40 Ligand is expressed as a homotrimer on platelets and activated T cells and B cells. It is upÂ-regulated following stimulation of basophils, eosinophils, fibroblasts, mast cells, monocytes, natural killer cells, vascular endothelial cells, and smooth muscle cells. CD40 Ligand binds and activates CD40, which is expressed on the surface of B cells, dendritic cells,macrophages, monocytes, platelets, endothelial cells, and epithelial cells. Monomeric, dimeric, and trimeric forms of soluble CD40 Ligand bind to oligomeric CD40 on cell membranes. CD40 ligation by CD40 Ligand promotes B cell activation and T cellÂdependent humoral responses. CD40 Ligand dysregulation on T cells and antigen presenting cells contributes to the immune deficiency associated with HIV infection and AIDS. It is also implicated in the pathology of multiple cardiovascular diseases including atherosclerosis, atherothrombosis, and restenosis.