

## Product datasheet for **TP727855**

### TNFRSF4 Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Human OX40 (C-mFc)
Species:	Human
Expression cDNA Clone or AA Sequence:	Leu29-Ala216
Tag:	C-mFc
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Note:	Recombinant Human OX40L Receptor is produced by our Mammalian expression system and the target gene encoding Leu29-Ala216 is expressed with a mFc tag at the C-terminus.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	12 months from date of despatch
Locus ID:	7293
UniProt ID:	<a href="#">P43489</a>
Synonyms:	Tumor necrosis factor receptor superfamily member 4;ACT35 antigen;OX40L receptor;TAX transcriptionally-activated glycoprotein 1 receptor;TNFRSF4;OX40;CD134;Txgp1
Summary:	OX40,also termed CD134 and TNFRSF4, is a T cell co-stimulatory molecule of the TNF receptor superfamily which plays a key role in the survival and homeostasis of effector and memory T cells. OX40 is expressed on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. The interaction between OX40 and OX40 ligand (OX40L) will occur when activated T cells bind to professional antigen-presenting cells (APCs). The T-cell functions, including cytokine production, expansion, and survival, are then enhanced by the OX40 costimulatory signals. OX40 signals are critical for controlling the function and differentiation of Foxp3+ regulatory T cells. OX40-OX40L interaction regulates T-cell tolerance, peripheral T-cell homeostasis, and T-cell-mediated inflammatory diseases.
Protein Families:	Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction


[View online »](#)