

## **Product datasheet for TP727882**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **MUC16 Human Recombinant Protein**

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant Human CA125 (C-6His)

Species: Human

**Expression cDNA Clone** 

or AA Sequence:

Gly12660-Met12923

Tag: C-His

**Buffer:** Lyophilized from a 0.2 um filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH7.5.

**Note:** Recombinant Human Mucin-16 is produced by our Mammalian expression system and the

target gene encoding Gly12660-Met12923 is expressed with a 6His tag at the C-terminus.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Stability:** 12 months from date of despatch

**Locus ID:** 94025 **UniProt ID:** 08WXI7

Synonyms: CA125 ovarian cancer antigen; CA125; CA-125; CA125MUC-16; FLJ14303; MUC16; mucin 16,

cell surface associated; mucin-16

Summary: MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I

transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. MUC16 is over-expressed by tumor cells including ovarian and mesothelial cancers. The transmembrane form can adhere to mesothelin in the peritoneum, facilitating metastasis of ovarian cancer to the peritoneal cavity. MUC16 also binds galectin-1 on immune cells and enhances its expression on tumor cells. MUC16-expressing tumors adhere to NK cells, down-regulate CD16 and suppress NK response, which may promote immune evasion. MUC16 is also cyclically expressed in the endometrium and may contribute to immune privilege during pregnancy. In the eye, MUC16 and other mucins protect the cornea and keep

it hydrated. It is altered on the conjunctival epithelium of patients with non-Sjogren dry eye

syndrome.

