

## **Product datasheet for TP727894**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Cynomolgus Recombinant Protein**

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant Cynomolgus Siglec-15 (C-6His)

Species: Cynomolgus
Expression cDNA Clone Phe20-Thr263

Expression cDNA Clone

or AA Sequence:

Tag: C-6His

**Buffer:** Lyophilized from a 0.2 um filtered solution of PBS,150mMNacl,0.3%chaps,5%Trehalose,

pH7.4.

**Note:** Recombinant Cynomolgus Sialic Acid-binding Ig-like lectin 15 is produced by our Mammalian

expression system and the target gene encoding Phe20-Thr263 is expressed with a 6His tag at

the C-terminus.

Stability: 12 months from date of despatch

Summary: Human Siglec-15 is a transmembrane glycoprotein in the Siglec family. Siglecs are type I

transmembrane proteins where the NH3+-terminus is in the extracellular space and the COOâ^'-terminus is cytosolic. Each Siglec contains an N-terminal V-type immunoglobulin domain (Ig domain) which acts as the binding receptor for sialic acid. These lectins are placed into the group of I-type lectins because the lectin domain is an immunoglobulin fold. All Siglecs are extended from the cell surface by C2-type Ig domains which have no binding activity. Siglecs differ in the number of these C2-type domains. Human Siglec-15 consists of a

244 amino acid (aa) extracellular domain (ECD) with two lg-like domains, a 21 aa

transmembrane segment, and a 44 aa cytoplasmic domain. Siglec-15 function is important

for osteoclast formation and TRANCE/RANK Ligand signaling in osteoclasts