

## Product datasheet for TP728024

## **BCMA (TNFRSF17) Human Recombinant Protein**

## **Product data:**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type:                            | Recombinant Proteins  |
|--|---|
| Description:                             | Biotinylated Human BCMA (C-Fc-Avi)  |
| Species:                                 | Human   |
| Expression cDNA Clone<br>or AA Sequence: | Met1-Ala54  |
| Tag:                                     | C-Fc-Avi  |
| Buffer:                                  | Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.   |
| Note:                                    | Biotinylated Recombinant Human B-cell Maturation Protein is produced by our Mammalian expression system and the target gene encoding Met1-Ala54 is expressed with a Fc, Avi tag at the C-terminus.  |
| Storage:                                 | Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.   |
| Stability:                               | 12 months from date of despatch   |
| Locus ID:                                | 608   |
| UniProt ID:                              | <u>Q02223</u>   |
| Synonyms:                                | Tumor necrosis factor receptor superfamily member 17;TNFRSF17;B-cell maturation protein;CD269   |
| Summary:                                 | B cell maturation protein (BCMA) is a type III membrane protein which belongs to the TNF receptor superfamily (TNFRSF17). BCMA contains one extracellular cysteine rich domain, as well as TACI. Human BCMA is a 184 amino acid (aa) protein consisting of a 54 aa extracellular domain, a 23 aa transmembrane domain, and a 107 aa intracellular domain. Mouse and human BCMA share 62% amino acid identity. BCMA is mainly expressed in immune organs and mature B cell lines. BCMA appears to be localized to the Golgi compartment. BCMA may play an important role in B cell development, function and regulation. |
| Protein Families:                        | Druggable Genome, Transmembrane   |
| Protein Pathways:                        | Cytokine-cytokine receptor interaction  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US