

Product datasheet for TP728044

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LAIR1 Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Biotinylated Human LAIR1 (C-Avi-6His)

Species: Human

Expression cDNA Clone

or AA Sequence:

UniProt ID:

Gln22-His163

Tag: C-Avi-6His

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Note: Biotinylated Recombinant Human Leukocyte-Associated Immunoglobulin-Like Receptor 1 is

produced by our Mammalian expression system and the target gene encoding Gln22-His163

is expressed with a AVI, 6His tag at the C-terminus.

Stability: 12 months from date of despatch

Q6GTX8

Locus ID: 3903

Summary: Leukocyte-Associated Immunoglobulin-Like Receptor 1 (LAIR1) is a single-pass type I

membrane protein. LAIR1 expressed on the majority of peripheral mononuclear cells, including natural killer (NK) cells, T-cells, B-cells, monocytes, and dendritic cells, highly in naive T-cells and B-cells. As an inhibitory receptor, LAIR1 plays a constitutive negative regulatory role on cytolytic function of natural killer (NK) cells, B-cells and T-cells. Activation by Tyr phosphorylation results in recruitment and activation of the phosphatases PTPN6 and PTPN11. It also reduces the increase of intracellular calcium evoked by B-cell receptor ligation. LAIR1 plays inhibitory role independently of SH2-containing phosphatases and modulates cytokine production in CD4+ T-cells. It down-regulates IL2 and IFNG production

production in B-cells as well as IL8, IL10 and TNF secretion. LAIR1 inhibits the differentiation of peripheral blood precursors towards dendritic cells. It also restrains proliferation and induces apoptosis in myeloid leukemia cell lines as well as prevents nuclear translocation of NF-kappa-B p65 subunit/RELA and phosphorylation of I-kappa-B alpha/CHUK in these cells.

while inducing secretion of transforming growth factor beta, also down-regulates IgG and IgE

