

## Product datasheet for **TP728144L**

### Recombinant Activin B, Human


#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant Activin B, Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGLECDGRTNLCCRQQFFIDFRLIGWNDWIIAPTGGYGN YCEGSCPAYLAGVPGSASSFHTAVVNQYRMR GLNPGTVNSCCIPTKLSTMSMLYFDDEYNIVKRDVPMIVEECGCA with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 13.75 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce hemoglobin expression in K562 cells. The ED <sub>50</sub> for this effect is <0.7 ng/mL. The specific activity of recombinant human Activin B is > 1.5 x 10 <sup>6</sup> IU/mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	<a href="#">P09529</a>
Synonyms:	Activin Beta B, INHBB; inhibin beta B chain


[View online »](#)

**Summary:**

Activin B is the cytokine of the TGF-beta family, which is a 13 kDa protein containing 115 amino acid residues. Activin B has the wide variety of biological function including stem cell differentiation, inflammation, bone remodeling and neural development. They also have a role in regulation of FSH secretion. Activin B induces the phosphorylation of Smad which binds the smad binding element.

**Product images:**


SDS- PAGE analysis of recombinant human Activin B