

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP728146L

Recombinant APRIL (A proliferation-inducing ligand), Human

Product data:

| Product Type: | Recombinant Proteins |
|---|--|
| Description: | Recombinant APRIL (A proliferation-inducing ligand), Human |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | MAVLTQKQKKQHSVLHLVPINATSKDDSDVTEVMWQPALRRGRGLQAQGYGVRIQDAGVYLLYSQVLF QDVTFTMGQVVSREGQGRQETLFRCIRSMPSHPDRAYNSCYSAGVFHLHQGDILSVIIPRARAKLNLSPH GTFLGFVKL with polyhistidine tag at the C-terminus. |
| Tag: | His Tag (C-term) |
| Predicted MW: | The protein has a calculated MW of 17.29 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis). |
| Purity: | >98% as determined by SDS-PAGE. |
| Buffer: | The protein was lyophilized from a 0.2 μm filtered solution containing 0.1% sarkosyl in 1X PBS, pH 8.0. |
| Dioactivity | Massured by its ability to induce call death in lurkat calls. The ED for this effect is 2.6.4.0 |
| Bioactivity: | Measured by its ability to induce cell death in Jurkat cells. The ED50 for this effect is 2.6-4.0 μ g/mL. |
| Endotoxin: | |
| - | μg/mL. |
| Endotoxin: | μg/mL. <0.1 EU per 1 μg of the protein by the LAL method. Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 μg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. |
| Endotoxin: Reconstitution Method: | μg/mL. <0.1 EU per 1 μg of the protein by the LAL method. Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 μg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. |
| Endotoxin: Reconstitution Method: Applications: | μg/mL. <0.1 EU per 1 μg of the protein by the LAL method. Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 μg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. Cell culture Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at - |
| Endotoxin: Reconstitution Method: Applications: Storage: | μg/mL. <0.1 EU per 1 μg of the protein by the LAL method. Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 μg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. Cell culture Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles. |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Summary: APRIL is a member of tumor necrosis factor ligand superfamily that expressed by stromal tissue, macrophages, and T cells. APRIL is a 27.4 kDa protein containing 250 residues, which plays a critical role in modulating tumor progression. Besides, APRIL has been demonstrated to involve in regulating B and T cell survival, proliferation and differentiation via binding with TNFRSF13B/TACI and TNFRSF17/BCMA.

Product images:



SDS- PAGE analysis of recombinant human APRIL

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US