

## Product datasheet for **TP728147S**

### Recombinant BAFF (B-cell activating factor), Human

#### Product data:


<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant BAFF (B-cell activating factor), Human
<b>Species:</b>	Human
<b>Expression Host:</b>	E. coli
<b>Expression cDNA Clone or AA Sequence:</b>	MAVQGP EETVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSAL EEEKENKILVKETGYFFIYGQVLYTDKT YAMGH LIQRKKVHVFGDELSLVT LFR CIQNPETLPNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDV TFFGALKLL with polyhistidine tag at the C-terminus.
<b>Tag:</b>	His Tag (C-term)
<b>Predicted MW:</b>	The protein has a calculated MW of 17.98 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
<b>Purity:</b>	>98% as determined by SDS-PAGE.
<b>Buffer:</b>	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
<b>Bioactivity:</b>	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED <sub>50</sub> for this effect is <0.5 ng/mL.
<b>Endotoxin:</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Reconstitution Method:</b>	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
<b>Applications:</b>	Cell culture
<b>Storage:</b>	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
<b>UniProt ID:</b>	<a href="#">Q9Y275</a>
<b>Synonyms:</b>	TNFSF13B, BLYS, CD257, DTL, TALL-1, TALL1, THANK, TNFSF20, TNLG7A, ZTNF4



[View online »](#)

**Summary:**

BAFF also known as BLYS, TALL-1 and TNFSF13B, which belongs to tumor necrosis factor family. BAFF is a 31.2 kDa type II transmembrane protein containing 285 residues that predominantly produced by myeloid cells. BAFF has been demonstrated to activate the survival of B-cells and the B-cell response by binding to BAFFR/BR3. Additionally, BAFF also takes part in regulating B and T cell function via forming two ligands-two receptors pathway through sharing TNFRSF13B/TACI and TNFRSF17/BCMA receptors with APRIL.

**Product images:**

SDS- PAGE analysis of recombinant human BAFF