

Product datasheet for **TP728157L**

Recombinant BMP-3 (Bone morphogenetic protein-3), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant BMP-3 (Bone morphogenetic protein-3), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MQWIEPRNCARRYLKVDFADIGWSEWIISPKSFDAYYCSGACQFPMPKSLKPSNHATIQSIVRAVGVVPGIPEPCCVPEKMSSLSILFFDENKNVVLKVYPNMTVESCACR with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 13.34 kDa. The protein migrates as 14 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>95% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.
Bioactivity:	Measure by its ability to inhibit BMP-2-induced alkaline phosphatase production by ATDC5 cells. The ED ₅₀ for this effect is < 10 µg/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P12645
Synonyms:	Osteogenin, BMP-3A



[View online »](#)

Summary:

Bone Morphogenetic Protein-3 (BMP-3) is an extracellular multifunctional signaling cytokine that is also a member of the TGF- β family. BMP-3 can bind with TGF- β receptor and participate in SMAD protein signal transduction through triggering pathway-restricted SMAD protein phosphorylation. The primary function of BMP-3 is osteoblast differentiation and induces bone formation.

Product images:

SDS- PAGE analysis of recombinant human BMP-3