

Product datasheet for TP728160

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant BMP-6 (Bone morphogenetic protein-6) Human

Product data:

Product Type: Recombinant Proteins

Description: Recombinant BMP-6 (Bone morphogenetic protein-6) Human

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MVSSASDYNSSELKTACRKHELYVSFQDLGWQDWIIAPKGYAANYCDGECSFPLNAHMNATNHAIVQTL VHLMNPEYVPKPCCAPTKLNAISVLYFDDNSNVILKKYRNMVVRACGCH with polyhistidine tag at the

C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 14.07 kDa. The protein migrates as 13 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 1×PBS, pH 8.0.

Bioactivity: Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED₅₀ for

this effect is <87 ng/mL.

Endotoxin: $< 0.1 \text{ EU per 1} \mu \text{g of the protein by the LAL method.}$

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 μ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P22004

Synonyms: VGR, VG-1-related protein






Summary:

Bone Morphogenetic Protein-6 (BMP-6) is an extracellular multifunctional cytokine that is also a member of the TGF- β family. BMP-6 can bind with the TGF- β receptor and triggers SMAD protein signal transduction. It can keep joint integrity and stability in adults and plays a vital role in regulating hepcidin to maintain iron ions in the body.

Product images:



SDS- PAGE analysis of recombinant human BMP-