

## Product datasheet for **TP728176M**

### Recombinant CXCL12 (24-88) (C-X-C motif chemokine 12), Human

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant CXCL12 (24-88) (C-X-C motif chemokine 12), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MVLSYRCPCRFFESHVARANVKHLKILNTPNCALQIVARLKNNNRQVCIDPKLKWIQEYLEKALN with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 8.55 kDa. The protein migrates as 10 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.1 M NaCl, pH 4.5.
Bioactivity:	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR4. The ED <sub>50</sub> for this effect is <0.5 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	<a href="#">P48061</a>
Synonyms:	IRH, PBSF, SCYB12, SDF1, TLSF, TPAR1



[View online »](#)

**Summary:**

C-X-C motif chemokine 12 (CXCL12) also named stromal cell-derived factor 1 (SDF-1), which is a chemokine of the intercrine alpha family. CXCL12 is a 19.5kDa protein containing 167 amino acid residues. The function of CXCL12 include the hematopoiesis, angiogenesis embryogenesis, inflammation and angiogenesis. CXCL12 activated by the IL-1, TNF or lipopolysaccharide and induce the immune response in leukocytes.

**Product images:**

SDS- PAGE analysis of recombinant human CXCL12