

Product datasheet for **TP728179S**

Recombinant CXCL3 (C-X-C motif chemokine 3), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant CXCL3 (C-X-C motif chemokine 3), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	ASVTELRQCQLQTLQGIHLKNIQSVNVRSPGPHCAQTEVIATLKNGKKACLNPASPMVQKIIKILNKGST N with polyhistidine tag at the N-terminus.
Tag:	His Tag (N-term)
Predicted MW:	The protein has a calculated MW of 8.67 kDa. The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED ₅₀ for this effect is <2 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P19876
Synonyms:	GRO-γ; MGSAY, MIP-2β, GRO3



[View online »](#)

Summary:

C-X-C motif chemokine 3 (CXCL3) also named Growth-regulated oncogene gamma (GRO γ), which is a chemokine of the intercrine alpha family. CXCL3 is a 8kDa protein containing 73 amino acid residues. During inflammation, CXCL3 is activated by the TNF and IL-1 which mediates the monocytes migration and adhesion by targeting the CXCR2. CXCL3 activates the JNK activity which induce the cell differentiation.

Product images:

SDS- PAGE analysis of recombinant human CXCL3