

## Product datasheet for **TP728190L**

### Recombinant FGF-11 isoform 1 (Fibroblast growth factor-11 isoform 1), Human

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant FGF-11 isoform 1 (Fibroblast growth factor-11 isoform 1), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MAALASSLIRQKREVREPGGSRPVSAQRRVCPRGTKSLCQKQLLILLSKVRLCGGRPARPDRGPEPQLKGI VTKLFCRQGFYLQANPDGSIQGTPEDTSSFTHFNLIPLVGLRVVTIQSAKLGHYMAMNAEGLLYSSPHFTAE CRFKECVFENYYVLYASALYRQRSGRAWYLGLDKEGQVMKGNRVKKTAAAHFLLPKLLEVAMYQEPSLH SVPEASPSSPPAP with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 25.81 kDa. The protein migrates as 30 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is <0.2 ng/mL.
Endotoxin:	<0.01 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	<a href="#">Q92914</a>
Synonyms:	FHF-3, FHF3



[View online »](#)

**Summary:**

The Fibroblast Growth Factors (FGF) family has two groups, secreted FGFs and intracellular FGFs (iFGFs). iFGFs are related to voltage-gated sodium (Nav) channels. Human FGF-11 isoform 1 is a 25 kDa cytokine with 225 amino acid residues.