

Product datasheet for **TP728200M**

Recombinant FGF-20 (Fibroblast growth factor-20), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant FGF-20 (Fibroblast growth factor-20), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MPLAEVGGFLGGLEGLGQQVGSFLLPPAGERPPLLGERSSAAERSARGGPGAAQLAHLHGILRRRQLYCR RTGFHLQILPDGSVQGTRQDHSLEFGILEFISVAVGLVLSIRGVDSGLYLG MNDKGEYSEKLTSECIFREQFE ENWYNTYSSNIYKHGDTGRRYFVALNKDGTPRDGARSKRHQKFTHFLPRPVDPERVPELYKDLLMYT with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 24.24 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is 1.3-3.2 ng/mL. The specific activity of recombinant human FGF-20 is > 2 x 10 ⁵ IU/mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q9NP95
Synonyms:	FGFK



[View online »](#)

Summary:

Fibroblast Growth Factors-20 (FGF-20) is a 23.5 kDa member of the fibroblast Growth Factors with 211 amino acid residues. FGF-20 is mainly expressed from microglial cells. FGF-20 can regulate central nervous development and function, regulate the survival of dopaminergic neurons via FGF/FGFR-dependent signaling pathway.

Product images:

SDS- PAGE analysis of recombinant human FGF-20