

Product datasheet for TP728207L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant FGF-6 (Fibroblast growth factor-6), Human

Product data:

Product Type: Recombinant Proteins

Description: Recombinant FGF-6 (Fibroblast growth factor-6), Human

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MGTRANNTLLDSRGWGTLLSRSRAGLAGEIAGVNWESGYLVGIKRQRRLYCNVGIGFHLQVLPDGRISGT HEENPYSLLEISTVERGVVSLFGVRSALFVAMNSKGRLYATPSFQEECKFRETLLPNNYNAYESDLYQGTYIA

LSKYGRVKRGSKVSPIMTVTHFLPRI with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 19.66 kDa. The protein migrates as 19 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate,

0.2 M NaCl, pH 3.5.

Bioactivity: Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <0.1 ng/mL.

The specific activity of recombinant human FGF-6 is $> 1 \times 10^7$ IU/mg.

Endotoxin: <0.1 EU per 1 μg of the protein by the LAL method.

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 μ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P10767

Synonyms: HBGF-6, HST-2






Summary:

Fibroblast Growth Factors-6 (FGF-6) is a 22.9 kDa member of the fibroblast Growth Factors with 208 amino acid residues. FGF-6 is an important role in the regulation of cell proliferation, cell differentiation, angiogenesis and myogenesis. In physiological function, FGF6 is required for normal muscle regeneration.

Product images:



SDS- PAGE analysis of recombinant human FGF-6