

Product datasheet for **TP728230**

Recombinant IGFBP7(IGF-binding protein 7) Human HEK293

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IGFBP7(IGF-binding protein 7) Human HEK293
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	A DNA sequence encoding Human IGFBP7 (#Q16270) (Ser27 – Leu282) was expressed with polyhistidine-SUMO tag at the N-terminus.
Tag:	His-SUMO Tag (N-term)
Predicted MW:	The protein has a calculated MW of 38.18 kDa. The protein migrates as 55 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>95% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Testing in process
Endotoxin:	<1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	Q16270
Synonyms:	IBP-7, IGFBP-7, IGFBP-rP1, MAC25, PSF(Prostacyclin-stimulating factor), TAF(Tumor-derived adhesion factor)



[View online »](#)

Summary:

IGFBP7, also known as mac25, prostacyclin-stimulating factor (PSF), tumor adhesion factor (TAF), and angiomodulin (AGM), is a secreted protein that belongs to the insulin-like growth factor (IGF)-binding protein (IGFBP) family. In contrast to the other family members, IGFBP7 binds IGFs through the N-terminal domain with low affinity. IGFBP7 is expressed in various human tissues, including the brain, liver, pancreas, and skeletal muscle, and is secreted into circulation. IGF-BP7 modulates many biological functions, such as protein synthesis, proliferation, anti-apoptosis, and cell survival. Moreover, it has been reported that IGFBP7 promotes cardiac senescence through IGF-1R/IRS/AKT-dependent suppression of FOXO3a, inhibiting DNA repair, subsequently the leading to the progression of Heart failure. Human and mouse IGFBP7 are highly homologous, sharing 94% a.a. sequence identity.

Product images:

SDS- PAGE analysis of recombinant human IGFBP7