

Product datasheet for **TP728236L**

Recombinant IL-11 (Interleukin-11), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-11 (Interleukin-11), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	PGPPPGPPRVSPDPRAELDSTVLLTRSLADTRQLAAQLRDKFPADGDHNLDSLPTLAMSAGALGALQLP GVLTRLRADLLSYLRHVQWLRRAGGSSSLKLEPELGTLQARLDRLRLQLLMSRLALPQPPDPAPPLA PPSSAWGGIRAAHAILGGLHLTLDWAVRGLLLKTRL with polyhistidine tag at the N-terminus.
Tag:	His Tag (N-term)
Predicted MW:	The protein has a calculated MW of 19.95 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce T11 cells proliferation. The ED ₅₀ for this effect is <0.2 ng/mL. The specific activity of recombinant human IL-11 is approximately >1 x 10 ⁷ IU/ mg.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P20809
Synonyms:	AGIF (Adipogenesis Inhibitory Factor)



[View online »](#)

Summary:

Interleukin 11 (IL-11) is a cytokine, encoded by the gp130 family genes. IL-11 initially thought to be important for hematopoiesis, notably for megakaryocyte maturation, and it facilitate the formation of higher order structures involving dimers of gp130: IL-11:IL-11 RA complexes.

Product images:

SDS- PAGE analysis of recombinant human IL-11