

Product datasheet for **TP728237S**

Recombinant IL-12 (p70) (Interleukin-12 p70), Human, HEK293

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant IL-12 (p70) (Interleukin-12 p70), Human, HEK293
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	IWELKKDVYVVELDWYPDAPGEMVLTCDTPEEDGITWTLQDQSEVLGSGKTLTIQVKEFGDAGQYTCHK GGEVLSHSLLLLHKKEDGIWSTDILKDQKEPKNKTLRCEAKNYSGRFTCWWLTTISTDLTFSVKSSRGSSD PQGVTCGAATLSAERVGRDNKEYEYSVECQEDSACPAAEESLPIEVMVDAVHKLKYENYSSFFIRDIKPD PKNLQLKPLKNSRQVEVSWEYPDTWSTPHSYFSLTFCVQVQGKSKREKDRVFTDKTSATVICRKNASISV RAQDRYSSSWSEWASVPCSGSTSGSGKPGSGEGSTKGRNLPVATPDPGMFPLHHSQNLRAVSNML QKARQTLFYPCTSEEDHEDITKDKTSTVEACLPLELTKNESCLNSRETSFITNGSCLASRKTSFMMALCLSS IYEDLKMYQVEFKTMNAKLLMDPKRQIFLDQNMLAVIDELMQALNFNSETVPQKSSLEEPDFYKTKIKLCI LLHAFRIRAVTIDRVMSYLNASHHHHHH with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 59.55 kDa. The protein migrates as 63-75 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.
Bioactivity:	Measure by its ability to induce IFN gamma secretion in PHA-activated human peripheral blood lymphocytes (PBMC). The ED ₅₀ for this effect is 0.05-0.2 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.




[View online »](#)

UniProt ID: [P29459](#)

Synonyms: Interleukin-12, NKSF, Cytotoxic Lymphocyte Maturation Factor (CLMF), TSF

Summary: Interleukin 12 p70 (IL-12 p70) is an interleukin which is major secreted from immune cells like dendritic cells, macrophages and neutrophils. IL-12 p70 is a 58.5 kDa protein containing 527 amino acid residues which is composed by two subunits (IL-12 p35 and IL-12 p40). IL-12 p70 induces the cytotoxic activity of immune cells like NK cells and T cells. IL-12 p70 binds its receptor (IL-12R β 1 and IL-12R β 2) that activates the Jak/STAT signaling pathway. IL-12 p70 also plays an important role in cancer model which inhibits the formation of new blood vessels through the production of IFN γ .

Product images:



SDS- PAGE analysis of recombinant human IL-12 p70