

Product datasheet for **TP728240S**

Recombinant IL-13 (Interleukin-13), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-13 (Interleukin-13), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGPVPPSTALRELIEELVNITQNQKAPLCNGSMVWSINLTAGMYCAALESINVSIGCSAIEKTQRMLSGFC PHKVSAGQFSSLHVRDTKIEVAQFVKDLLLHLKFLFREGRFN with polyhistidine tag at the C-terminus
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 13.28 kDa. The protein migrates as 15 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>95% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce TF-1 cells proliferation. The ED ₅₀ for this effect is <0.8 ng/mL. The specific activity of recombinant human IL-13 is approximately >1 x10 ⁶ IU/ mg.
Endotoxin:	<0.01 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P35225
Synonyms:	NC30



[View online »](#)

Summary:

Interleukin 13 (IL-13) with a molecular mass of 10 kDa cytokine, it secreted by T helper type 2 (Th2) cells, CD4 cells, natural killer T cell, mast cells, basophils, eosinophils and nuocytes. It is homologous to IL-4 and shares many of its biologic activities on mononuclear phagocytic cells, endothelial cells, epithelial cells, and B cells.

Product images:

SDS- PAGE analysis of recombinant human IL-13