

## Product datasheet for TP728242L

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Recombinant IL-17A (Interleukin-17A), Human

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant IL-17A (Interleukin-17A), Human

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MIVKAGITIPRNPGCPNSEDKNFPRTVMVNLNIHNRNTNTNPKRSSDYYNRSTSPWNLHRNEDPERYPSV IWEAKCRHLGCINADGNVDYHMNSVPIQQEILVLRREPPHCPNSFRLEKILVSVGCTCVTPIVHHVA with

polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

**Predicted MW:** The protein has a calculated MW of 16.47 kDa. The protein migrates as 17 kDa under

reducing condition (SDS-PAGE analysis).

**Purity:** >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.

**Bioactivity:** Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED $_{50}$  for this effect is <6 ng/mL.

**Endotoxin:** <0.01 EU per 1 μg of the protein by the LAL method.

**Reconstitution Method:** Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile  $H_2O$  to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**Applications:** Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: Q16552
Synonyms: CTLA-8

Summary: Interleukin 17A (IL-17A) predicts a molecular mass of 35 kDa, encoded IL17A gene is a


proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities

of NF-kappa B and mitogen-activated protein kinases.





# **Product images:**



SDS- PAGE analysis of recombinant human IL-17A