

Product datasheet for TP728248

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant IL-20 (Interleukin-20), Human

Product data:

Product Type: Recombinant Proteins

Description: Recombinant IL-20 (Interleukin-20), Human

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MLKTLNLGSCVIATNLQEIRNGFSEIRGSVQAKDGNIDIRILRRTESLQDTKPANRCCLLRHLLRLYLDRVFK

NYQTPDHYTLRKISSLANSFLTIKKDLRLCHAHMTCHCGEEAMKKYSQILSHFEKLEPQAAVVKALGELDIL

LQWMEETE with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 18.47 kDa. The protein migrates as 16 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0.

Bioactivity: Measure by its ability to chemoattract BaF3 cells transfected with human IL-20R alpha and IL-

20R beta. The ED₅₀ for this effect is <0.2 ng/mL.

Endotoxin: $< 0.1 \text{ EU per 1} \mu \text{g of the protein by the LAL method.}$

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

lyophilized protein in sterile H_2O to a concentration not less than 100 $\mu g/mL$ and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: Q9NYY1

Synonyms: Cytokine ZCYTO10






Summary:

Interleukin 20 (IL-20) predicts a molecular mass of 17.6 kDa. The IL-20 subfamily of cytokines is comprised of IL-19, IL-20, IL-22, IL-24, and IL-26. It regulates of differentiation of keratinocytes during inflammation, the expansion of multipotential hematopoietic progenitor cells.

Product images:



SDS- PAGE analysis of recombinant human IL-20