

Product datasheet for **TP728258M**

Recombinant IL-29 (Interleukin-29), Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-29 (Interleukin-29), Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MPTSKPTTTGKGCHIGRFKSLSPQELASFKKARDALEESLKLKNWSCSSPVFPGNWDLRLLQVRERPVALEAELALTLKVLEAAAGPALEDVLDQPLHLHILSQLQACIQPQPTAGPRPRGRLHHWLHRLQEAPKKESA GCLEASVTFNLFRLLTRDLKYVADGNLCLRTSTHPEST with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 20.70 kDa. The protein migrates as 22 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce IL-8 secretion in HuH7 cells. The ED ₅₀ for this effect is <6 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	<u>Q8IU54</u>
Synonyms:	IFN-λ1



[View online »](#)

Summary:

Interleukin 29 (IL-29) is a cytokine, predicts a molecular mass of 21.9 kDa. It belongs to type III interferons group, also termed interferons λ (IFN- λ). Its induction of STAT3-STAT5 has also been displayed, albeit to a lesser degree. The STAT1 /STAT2 signaling cascade transpires as follows: once tyrosine residues on STAT1 and STAT2 are phosphorylated, these proteins dimerize and are subsequently transported to the nucleus.

Product images:

SDS- PAGE analysis of recombinant human IL-29