

Product datasheet for TP728270M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Recombinant IL-5 (Interleukin-5), Human

Product data:

Product Type: Recombinant Proteins

Description: Recombinant IL-5 (Interleukin-5), Human

Species: Human **Expression Host:** E. coli

Expression cDNA Clone

MIPTEIPTSALVKETLALLSTHRTLLIANETLRIPVPVHKNHQLCTEEIFQGIGTLESQTVQGGTVERLFKNLSL or AA Sequence: IKKYIDGQKKKCGEERRRVNQFLDYLQEFLGVMNTEWIIES with polyhistidine tag at the C-terminus.

Tag: His Tag (C-term)

Predicted MW: The protein has a calculated MW of 14.09 kDa. The protein migrates as 14 kDa under

reducing condition (SDS-PAGE analysis).

Purity: >98% as determined by SDS-PAGE.

Buffer: The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.

Bioactivity: Measure by its ability to induce TF-1 cells proliferation. The ED₅₀ for this effect is <0.3 ng/mL.

Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.

Reconstitution Method: Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the

> lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Applications: Cell culture

Storage: Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to

> 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -

20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

UniProt ID: P05113

Synonyms: EDF, BCDFII, TRF

Summary: IL-5 is a 45-60 kDa homodimer cytokine that functions principally in the eosinophil arm of the


Th2 response. It is a member of the helical bundle family of cytokines, along with IL-3, IL-4,

and granulocyte-macrophage colony-stimulating factor (GM-CSF).





Product images:



SDS- PAGE analysis of recombinant human IL-5