

Product datasheet for **TP728273**

Recombinant IL-8 (aa28-99) (Interleukin-8) \square Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant IL-8 (aa28-99) (Interleukin-8) \square Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MSAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSDGRELCLDPKENWVQRWVEKFLKRAENS with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 9.32 kDa. The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to chemoattract PBMC. The ED ₅₀ for this effect is <2 ng/mL.
Endotoxin:	<0.1 EU per 1 μ g of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 μ g/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	P10145
Synonyms:	Monocyte-Derived Neutrophil Chemotactic Factor (MDNCF), Neutrophil Activating Factor (NAF), CXCL8, C-X-C motif chemokine ligand 8



[View online »](#)

Summary:

Human IL-8 (aa 28-99) predicts a molecular mass of 8.5 kDa and is the 72 amino acid residue form Ser 28 to Ser 99 of mature human IL-8. This recombinant protein is the major form secreted by macrophages. IL-8 is a chemokine produced by macrophages and other cell types such as epithelial cells, airway smooth muscle cells and endothelial cells.

Product images:

SDS- PAGE analysis of recombinant human IL-8