

Product datasheet for **TP728276M**

Recombinant LIGHT, Human

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant LIGHT, Human
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MRLGEMVTRLPDGPAGSWEQLIQERRSHEVNPA AHLTGANSSLTGSGGPLLWETQLGLAFLRGLSYHDG ALVVTKAGYYYIYSKVQLGGVGCPLGLASTITHGLYKRTPRYPEELELLVSQQSPCGRATSSSRVWWDSSFL GGVVHLEAGEKVVVRVLDERLRLRDGTRS YFGAFMV with polyhistidine tag at the C-terminus.
Tag:	His Tag (C-term)
Predicted MW:	The protein has a calculated MW of 20.30 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Purity:	>98% as determined by SDS-PAGE.
Buffer:	The protein was lyophilized from a 0.2 µm filtered solution containing 0.1% sarkosyl in 1X PBS, pH 8.0.
Bioactivity:	Measure by its ability to induce cytotoxicity in HT-29 cells in the presence of IFN-gamma. The ED ₅₀ for this effect is < 10 ng/mL.
Endotoxin:	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution Method:	Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
Applications:	Cell culture
Storage:	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
UniProt ID:	O43557.2
Synonyms:	TNFSF14, HVEM-L, CD258, LTg



[View online »](#)

Summary:

LIGHT, also known as TNFSF14 is a member of the TNF superfamily that produced by multiple immune cells such as activated T cells and immature dendritic cells (DCs). LIGHT is a 29 kDa type II transmembrane protein, which serves as a ligand for both lymphotoxin β receptor (LT β R) and TNFSF signaling receptors (TNFRSF14/HVEM). Besides, LIGHT can amplify NF- κ B signaling pathway in T cells when presenting anti-CD3 antibody treatment. Additionally, LIGHT can stimulate T cell proliferation and IFN-expression via interacting with TNFRS13 /HVEM.

Product images:

SDS- PAGE analysis of recombinant human LIGHT